

# GeoFile 2-2023 ReadMe

## Manitoba till-matrix geochemistry compilation: silt plus clay (<63 µm) size-fraction by inductively coupled plasma–mass spectrometry after an aqua-regia or modified aqua-regia digestion



by M.S. Gauthier  
Manitoba Geological Survey  
Winnipeg, 2023

---

© King's Printer for Manitoba, 2023

Every possible effort is made to ensure the accuracy of the information contained in this report, but Manitoba Economic Development, Investment, Trade and Natural Resources does not assume any liability for errors that may occur. Source references are included in the report and users should verify critical information.

Any third party digital data and software accompanying this publication are supplied on the understanding that they are for the sole use of the licensee, and will not be redistributed in any form, in whole or in part. Any references to proprietary software in the documentation and/or any use of proprietary data formats in this release do not constitute endorsement by Manitoba Economic Development, Investment, Trade and Natural Resources of any manufacturer's product.

When using information from this publication in other publications or presentations, due acknowledgment should be given to the Manitoba Geological Survey. The following reference format is recommended:

Gauthier, M.S. 2023: Manitoba till-matrix geochemistry compilation: silt plus clay (<63 µm) size-fraction by inductively coupled plasma–mass spectrometry after an aqua-regia or modified aqua-regia digestion; Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, GeoFile 2-2023, Microsoft® Excel® file.

### Published by:

Manitoba Economic Development, Investment, Trade and Natural Resources  
Manitoba Geological Survey  
360–1395 Ellice Avenue  
Winnipeg, Manitoba  
R3G 3P2 Canada

Telephone: 1-800-223-5215 (General Enquiry)

204-945-6569 (Publication Sales)

Fax: 204-945-8427

Email: [minesinfo@gov.mb.ca](mailto:minesinfo@gov.mb.ca)

Website: [manitoba.ca/minerals](http://manitoba.ca/minerals)

This publication supersedes GeoFile 2-2022.

This publication is available to download free of charge at [manitoba.ca/minerals](http://manitoba.ca/minerals)

## Abstract

This GeoFile provides a digital dataset for till-geochemistry surveys carried out in Manitoba, where the silt plus clay (<63 µm) size-fraction of the till matrix was analyzed by inductively coupled plasma–mass spectrometry after an aqua-regia or modified aqua-regia digestion. This compilation of 35 projects includes 3554 till samples, and will be updated annually or bi-annually. This data can be brought into GIS software, and integrated with other geoscience data, to generate new exploration targets and design follow-up exploration programs.

## Résumé

Ce géodossier offre un jeu de données numériques pour les relevés de géochimie du till effectués au Manitoba, pour lesquels la classe granulométrique du limon et de l'argile (<63 µm) de la matrice de till a été analysée au moyen d'une spectrométrie de masse à plasma à couplage inductif (ICP-MS) après digestion à l'eau régale ou à l'eau régale modifiée. Cette compilation de 35 projets comprend 3 554 échantillons de till et sera mise à jour une fois par an ou une fois tous les deux ans. Ces données peuvent être téléchargées dans un logiciel SIG et intégrées à d'autres données géoscientifiques, afin de générer de nouvelles cibles d'exploration et de concevoir des programmes d'exploration de suivi.

---

## DIGITAL DATA

Zip file geofile2.zip contains the following content:

- GeoFile\_2-2023\_ReadMe.pdf (this file)
- GeoFile\_2-2023.xlsx:
  - Table 1: Detection limits.
  - Table 2: Till-matrix ICP data on the silt plus clay (<63 µm) size-fraction.
  - Table 3: Summary statistics for the till-matrix ICP data on the silt plus clay (<63 µm) size-fraction.
  - Table 4: References.

## Introduction

This GeoFile captures till-matrix geochemistry data collected from surveys carried out in Manitoba since the 1990s. These surveys generally combine surficial mapping, paleo ice-flow mapping and sampling of till to be analyzed for geochemistry. Publication of the till-geochemistry data from these surveys have been successful in generating exploration activity, as well as providing ‘background’ baseline values.

Moving forward in a digital age, all till-geochemistry data is compiled into databases that will allow users to quickly view, compile and interact with the data from different regions of Manitoba. This data will enable users to more quickly identify when an element concentration is atypical for an area.

This GeoFile includes data from 3554 till samples collected between 1991 and 2023, where the silt plus clay (<63 µm) size-fraction of the till matrix was analyzed by inductively coupled plasma–mass spectrometry (ICP-MS) after an aqua-regia or modified aqua-regia digestion (Figure 1). This database will be updated as new data is released.

## Methods

### Updates

Data from Manitoba Geological Survey (MGS) fieldwork was added for northern Manitoba (Gauthier and Hodder 2022; Hodder and Gauthier 2022a, b; Gauthier and Hodder 2023b–d) and southeastern Manitoba (parts of 62H2, 7; Gauthier and Hodder, 2023a); as well as company data given to the MGS from the Snow Lake area (Assessment File 63K231070, Manitoba Economic Development, Investment, Trade and Natural Resources, Winnipeg). Two hundred far-northern Manitoba samples were re-analyzed, and the new data is recorded herein (Gauthier and Godbout, 2023). Coordinates were updated from the incorrect NAD27 to the correct NAD83 for Dredge and McMartin (2007), Thorleifson and Matile (1993) and Thorleifson et al. (2009). In addition, notes were added on the stratigraphy for southeastern Manitoba rotosonic drillcores (NTS 62H7; from Thorleifson and Matile, 1993), based on information in Matile et al. (2023).

### Collection methods

Till samples were collected from road cuts, borrow pits, ditches, natural exposures, hand-dug holes, Dutch-auger holes and boreholes across Manitoba. Wherever possible, till samples were collected from the C horizon in order to minimize potential weathering effects. To learn more about the characteristics of individual till samples, the reader is encouraged to view the original publication.

Till-geochemical surveys in Manitoba were first compiled to produce a till geochemistry index map. Following that, the data itself was manually compiled and separated according to size-fraction and analytical method. This compilation includes all data where the matrix (<63 µm size-fraction) was analyzed by ICP-MS

after an aqua-regia or modified aqua-regia digestion. There are also till compilations for analyses by instrumental neutron activation (Gauthier, 2020a), the clay size-fraction (Gauthier, 2020b), visible gold in the heavy mineral fraction (Gauthier, 2023b) and kimberlite-indicator minerals (Keller, 2019).


Data captured includes all data immediately relevant to the till sample. This includes publication number, laboratory used, project name, spatial coordinates, depth of sample and other important information. Depth of sample was transcribed from the original publications, and includes both depth ranges (e.g., 0.7–1.2 m) and single digits (e.g., 1.2 m). MGS project numbers are only assigned to some projects as this is a newer internal initiative designed to better track projects from year to year. The user should note that the compilation includes samples taken at depth, in some cases by drilling and in others accessed by natural river sections.

### Sample location

Sample locations are provided for each till sample. Technically, Manitoba crosses three UTM zones (14 to 16). For ease of display in GIS, all data has been re-projected into zone 14. Hence, all coordinates herein are reported as UTM zone 14, NAD83. Some older samples may be misplaced by as much as 200 m, as it is unknown when recording methods switched from NAD27 to NAD83. While coordinates were compiled from the original reports, some projects were pre-GPS and the locations were digitized from hand-drawn field maps. Again, the coordinates of these older till samples are to be used as a guide instead of a precise location. This is why Table 2 includes the column ‘Year\_sampled’ instead of the publication year.

### Analytical methods

Till samples were prepared for geochemistry analysis at the Saskatchewan Research Council Geoanalytical Laboratories (Saskatoon, Saskatchewan), the Geological Survey of Canada’s Sedimentary Laboratory (Ottawa, Ontario), the MGS Midland Sample and Core Library (Winnipeg, Manitoba), and perhaps other undisclosed labs. This compilation includes data from 35 Manitoba till-sampling projects where the matrix (<63 µm size-fraction) was analyzed by ICP-MS after an aqua-regia or modified aqua-regia digestion between 1991 and 2023. Data from five projects (Henderson, 1994; McClenaghan et al., 2009; Hodder, 2018; Gauthier, 2019; Gauthier and Hodder, 2020) where Au, Pd and/or Pt was analyzed by ICP-MS after fire assay is also included (FA\_ICP; Table 2). The processing and analytical methods used vary according to the date of the survey. Earlier surveys may have been analyzed for a restricted suite of elements, or at different detection limits (Table 1). Samples from the 35 projects were analysed at various labs over time, including the Saskatchewan Research Council Geoanalytical Laboratories, Acme Analytical Laboratories Ltd. (now Bureau Veritas Commodities Canada Ltd.; Vancouver, British Columbia), Activation Laboratories Ltd. (Ancaster, Ontario) and ALS Chemex (now ALS-Geochemistry; Vancouver, British



**Figure 1:** Till-sample locations where the silt plus clay (<63  $\mu\text{m}$ ) size-fraction of the matrix was analyzed by inductively coupled plasma–mass spectrometry after an aqua-regia or modified aqua-regia digestion in Manitoba.

Columbia). The full list of projects included, and elements analyzed, can be found in Table 4.

### **Compilation methods**

Included in this digital database is data from 3554 till samples collected from 35 different projects. No effort is made to re-analyze, level or otherwise standardize these values. Some studies reported Ag, Au, Hg and Re using different measurements—all have been reported as ppb in Table 2. Concentrations of P, reported as ppm in three studies (Table 3), have been converted to percent in Table 2.

Values below the detection limit are reported as the negative of the detection limit. This is because the detection limits for most elements vary in orders of magnitude over time (Table 1). The reader should assess the data accordingly.

## **Preliminary results**

### **Summary statistics**

The summary statistics for all reported elements are depicted in Table 3. Nine elements where the 90<sup>th</sup> percentile of values are at or below detection limit are shown in grey. Most of these elements are not useful for exploration purposes using ICP after aqua-regia or modified aqua-regia. Given that geology has spatial patterns, all data of interest should be analyzed spatially as well (Grunsky, 2010). For example, background values of iron will be lower for tills derived from carbonate bedrock than from granitoid bedrock.

### **Carbonate till distribution**

A significant portion of the till in Manitoba is calcareous (Figure 2). This carbonate has two sources: Paleozoic bedrock within the Hudson Bay Basin in the far northeast, and with the Western Canada Sedimentary Basin (WCSB) in the south (Wheeler et al., 1996). The net carbonate-dispersal pattern within the till is complex (Figure 2), and generally decreases in concentration to the west, southwest, and south of Hudson Bay. Then, the concentrations increase drastically within tills south of Flin Flon and Snow Lake, reflecting quick entrainment of calcareous detritus from the WCSB. Within this larger pattern, however, the calcareous surface tills locally contain a range of carbonate concentrations that relate to overprinting (dilution and/or reworking) and inheritance (preservation) during till transportation and deposition (e.g., Trommelen et al., 2013; Trommelen and Ross, 2014; Gauthier et al., 2019).

### **Prospective and background concentrations**

Because carbonate rocks are less resistant than most Precambrian shield rocks, they can mask, or dilute, the ‘signature’ of elements important to exploration. Within areas of Manitoba covered by calcareous till, ‘low’ concentrations of desired elements may still be more prospective than the same concentration within noncalcareous till. The total car-

bonate (wt. %) concentration is reported for most samples within Table 2. The entire dataset of Manitoba tills is plotted in Figure 3 for As, Cr, Cu, Ni and Zn analyzed by ICP-MS after an aqua-regia or modified aqua-regia digestion. For As and Ni, higher concentrations tend to occur when the total carbonate concentration of Manitoba tills is low. For the entire dataset, ‘low’ could be defined as 5% for arsenic (Figure 3a), and 35% for nickel (Figure 3d). ‘Elevated values’ of metals occurring at high carbonate concentration, such as the 30 ppm As measurement at 67 wt. % CO<sub>3</sub>, could require follow-up sampling when conducting drift exploration. For Cr, Cu and especially Zn, there is a negative correlation to the total carbonate concentration of Manitoba tills (Figure 3b, c, e). These are necessarily broad general statements, and there are anomalies within each graph. The reason ‘why’ a particular relationship occurs would depend on what bedrock the till is overlying, what bedrock types the till is sourced from, and what other materials may have been incorporated into the till (glaciolacustrine, glaciomarine, nonglacial sediment types, etc). In general, calcareous values should be noted and different populations should not necessarily be treated as one dataset.

### **Supporting data**

The original files for each project can be found through the Bibliography of Manitoba Geology and Resource Centre Catalogue (Manitoba Geological Survey, 2021a). To help with analysis, the following data is also available:


- Manitoba carbonate dispersal analyses in till (Gauthier, 2023a)
- index of surficial geology maps in Manitoba (Manitoba Geological Survey, 2021c)
- compiled surficial materials maps (Manitoba Geological Survey, 2021b)
- digital compilation of surficial point and line features, including ice-flow data (striations, streamlined landforms) and bedrock outcrop locations (Gauthier et al., 2022)
- the current understanding of ice-flow history in northeastern Manitoba (Gauthier et al., 2019)

## **References**

- Dredge, L.A. and McMartin, I. 2007: Geochemical reanalysis of archived till samples from northernmost Manitoba; Geological Survey of Canada, Open File 5546, 95 p.
- Gauthier, M.S. 2019: Till composition, stratigraphy, ice-flow indicator data and glacial history of the North Knife River–Churchill River region, Manitoba (parts of NTS 54L, 64I); Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Geoscientific Paper GP2019-1, 41 p., URL <<https://manitoba.ca/iem/info/libmin/GP2019-1.zip>> [September 2021].
- Gauthier, M.S. 2020a: Manitoba till-matrix geochemistry compilation 1: silt plus clay (<63 µm) size-fraction by instrumental neutron activation analysis; Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Open File OF2020-2, 6 p., URL <<https://manitoba.ca/iem/info/libmin/OF2020-2.zip>> [September 2021].







**Figure 3:** Plots of arsenic (a), chromium (b), copper (c), nickel (d) and zinc (e) concentration versus total carbonate concentration—for the entire dataset—show that the concentration of metals varies with the concentration of carbonate in the tills in Manitoba.

Gauthier, M.S. 2020b: Manitoba till-matrix geochemistry compilation 3: clay (<2 µm) size-fraction by atomic absorption spectrometry or inductively coupled plasma–emission spectrometry after aqua-regia digestion; Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Open File OF2020-5, 5 p., URL <<https://manitoba.ca/iem/info/libmin/OF2020-5.zip>> [September 2021].

Gauthier, M.S. 2023a: Manitoba till-matrix geochemistry compilation: total carbonate of the silt plus clay (<63 µm) size-fraction; Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, GeoFile 7-2023, Microsoft® Excel® file.

Gauthier, M.S. 2023b: Manitoba till-matrix geochemistry compilation: visible gold grains in the heavy mineral (<2 mm) size-fraction; Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, GeoFile 8-2023, Microsoft® Excel® file.

Gauthier, M.S. and Godbout, P.-M. 2023: Till-matrix geochemistry re-analyses data from far northeastern and far northwestern Manitoba (parts of NTS 54E, F, L, 64J, K, N, O); Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, Data Repository Item DRI2023007, Microsoft® Excel® file.

- Gauthier, M.S. and Hodder, T.J. 2020: Till-matrix geochemistry data from the Gillam area, northeast Manitoba; Manitoba Agriculture and Resource Development, Manitoba Geological Survey, Data Repository Item DRI2020005, Microsoft® Excel® file, URL <<https://manitoba.ca/iem/info/libmin/DRI2020005.xlsx>> [September 2021].
- Gauthier, M.S. and Hodder, T.J. 2022: Till-matrix geochemistry data from the Gillam area, northeastern Manitoba: additional 2021 data (NTS 54D7, 9, 11); Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, Data Repository Item DRI2022004, Microsoft® Excel® file.
- Gauthier, M.S. and Hodder, T.J. 2023a: Quaternary site data, till composition and ice-flow indicators in the Roseau River area, southeastern Manitoba (parts of NTS 62H2, 7); Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, Open File OF2023-3, 10 p., 8 appendices.
- Gauthier, M.S. and Hodder, T.J. 2023b: Till geochemistry and heavy mineral analyses (gold, MMSIM®, visual KIM) from three sections near the confluence of the Hayes and Gods rivers, northeastern Manitoba (part of NTS 54C7); Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, Data Repository Item DRI2023008, Microsoft® Excel® file.
- Gauthier, M.S. and Hodder, T.J. 2023c: Till geochemistry and heavy mineral analyses (gold, MMSIM®, visual KIM) of the western Fox River greenstone belt area, northeastern Manitoba (NTS 53M15, 16, parts of 53N13, 54C4, 54D1): year 2; Manitoba Economic Development, Investment, Trade and Natural Resources, Manitoba Geological Survey, Data Repository Item DRI2023015, Microsoft® Excel® file.
- Gauthier, M.S. and Hodder, T.J. 2023d: Till-matrix geochemistry data from the Gillam area, northeastern Manitoba: additional 2022 data (NTS 54D5–8, 64A2); Manitoba Economic Development, Investment and Trade, Manitoba Geological Survey, Data Repository Item DRI2023004, Microsoft® Excel® file.
- Gauthier, M.S., Santucci, A. and Keller, G.R. 2022: Digital compilation of surficial point and line features for Manitoba, including ice-flow data; Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, GeoFile 1-2022, 5 p., URL <<https://manitoba.ca/iem/info/libmin/geofile1.zip>> [February 2022].
- Gauthier, M.S., Hodder, T.J., Ross, M., Kelley, S.E., Rochester, A. and McCausland, P. 2019: The subglacial mosaic of the Laurentide Ice Sheet; a study of the interior region of southwestern Hudson Bay; *Quaternary Science Reviews*, v. 214, p. 1–27, URL <<https://doi.org/10.1016/j.quascirev.2019.04.015>>.
- Grunksy, E.C. 2010: The interpretation of geochemical survey data; *Geochemistry: Exploration, Environment, Analysis*, v. 10, no. 1, p. 27–74, URL <<https://doi.org/10.1144/1467-7873/09-210>>.
- Henderson, P.J. 1994: Surficial geology and drift composition of the Bissett-English Brook area, Rice Lake Greenstone Belt, southeastern Manitoba; Geological Survey of Canada, Open File 2910, 189 p., URL <<https://doi.org/10.4095/194153>>.
- Hodder, T.J. 2018: Ice-flow history and till composition of the Southern Indian Lake area, north-central Manitoba (parts of NTS 64G1, 2, 7–10, 64B15); Manitoba Growth, Enterprise and Trade, Manitoba Geological Survey, Open File OF2018-1, 21 p., URL <<https://manitoba.ca/iem/info/libmin/OF2018-1.zip>> [September 2021].
- Hodder, T.J. and Gauthier, M.S. 2022a: Till-matrix geochemistry data from the Churchill–Little Churchill rivers area, northeastern Manitoba (part of NTS 54E); Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, Data Repository Item DRI2022005, Microsoft® Excel® file.
- Hodder, T.J. and Gauthier, M.S. 2022b: Till-matrix geochemistry data from the Machichi–Kettle rivers area, far northeastern Manitoba (parts of NTS 54A–C); Manitoba Natural Resources and Northern Development, Manitoba Geological Survey, Data Repository Item DRI2022006, Microsoft® Excel® file.
- Keller, G.R. 2019: Manitoba Kimberlite Indicator Mineral Database (version 3.2); Manitoba Growth, Enterprise and Trade, Manitoba Geological Survey, zipped Microsoft® Access® 2016 database, URL <[https://manitoba.ca/iem/geo/diamonds/MBKIMDB\\_32.zip](https://manitoba.ca/iem/geo/diamonds/MBKIMDB_32.zip)> [September 2021].
- Manitoba Department of Mines, Natural Resources and Environment 1979: Geological map of Manitoba; Manitoba Department of Mines, Natural Resources and Environment, Mineral Resources Division, Geological Report GR79-2, scale 1:1 000 000.
- Manitoba Geological Survey 2021a: Introduction to the Bibliography of Manitoba Geology and Resource Centre catalogue (BMG); Manitoba Agriculture and Resource Development, Manitoba Geological Survey, URL <<https://manitoba.ca/iem/info/library/bmgintro.html>> [September 2021].
- Manitoba Geological Survey 2021b: Surficial geology compilation map series (SGCMS); Manitoba Agriculture and Resource Development, Manitoba Geological Survey, URL <<https://manitoba.ca/iem/geo/gis/surfgeomap.html>> [September 2021].
- Manitoba Geological Survey 2021c: Surficial geology map index; Manitoba Agriculture and Resource Development, Manitoba Geological Survey; URL <[https://manitoba.ca/iem/geo/surficial/sg\\_gf.html](https://manitoba.ca/iem/geo/surficial/sg_gf.html)> [September 2021].
- Matile, G.L.D., Thorleifson H. and Gauthier, M.S. 2023: Rotosonic borehole stratigraphy, southeastern Manitoba (parts of NTS 52E, 62H); Manitoba Economic Development, Investment and Trade, Manitoba Geological Survey, Data Repository Item DRI2023005, Microsoft® Excel® file, photos and PDFs.
- McClenaghan, M.B., Matile, G.L.D., Layton-Matthews, D. and Pyne, M. 2009: Till geochemical signatures of magmatic Ni-Cu deposits and regional till geochemistry, Thompson Nickel belt, Manitoba; Geological Survey of Canada, Open File 6005, 121 p., URL <<https://doi.org/10.4095/247412>>.
- Thorleifson, L.H. and Matile, G. 1993: Till geochemistry and indicator mineral reconnaissance of southeastern Manitoba; Geological Survey of Canada, Open File 2750, 2 diskettes.
- Thorleifson, L.H., Matile, G.L.D., Keller, G.R. and Hauck, S.A. 2009: Till geochemical and indicator mineral reconnaissance of southeastern Manitoba (west half of NTS 52E and 52L and all of 62H and 62I): final results; Manitoba Science, Technology, Energy and Mines, Manitoba Geological Survey, Open File OF2009-13, 5 p.
- Trommelen, M.S. and Ross, M. 2014: Distribution and type of sticky spots at the centre of a deglacial streamlined lobe in northeastern Manitoba, Canada; *Boreas*, v. 43, no. 3, p. 557–576, URL <<https://doi.org/10.1111/bor.12064>>.
- Trommelen, M.S., Ross, M. and Campbell, J.E. 2013: Inherited clast dispersal patterns: implications for paleoglaciology of the SE Keewatin Sector of the Laurentide Ice Sheet; *Boreas*, v. 42, no. 3, p. 693–713, URL <<https://doi.org/10.1111/j.1502-3885.2012.00308.x>>.
- Wheeler, J.O., Hoffman, P.F., Card, K.D., Davidson, A., Sanford, B.V., Okulich, A.V. and Roest, W.R. 1996: Geological map of Canada; Geological Survey of Canada, Map 1860A, 3 sheets, scale 1:5 000 000, URL <<https://doi.org/10.4095/208175>>.