

490 Hargrave Street Winnipeg, Manitoba R3A OX7 CANADA 490, rue Hargrave Winnipeg (Manitoba) R3A 0X7 CANADA

www.wrha.mb.ca

www.wrha.mb.ca

November 8, 2020

Transcona Collegiate 1305 Winona St Winnipeg, MB R2C 2P9

Dear Parent/Guardian of Students at Transcona Collegiate,

Manitoba public health officials advised Transcona Collegiate today of a confirmed case of COVID-19 present in the school on October 27-29, 2020 when the individual may have been infectious. An investigation is ongoing to determine if there is a link between this case and previous cases.

Transcona Collegiate is working closely with public health officials and following their recommendations. Based on the public health investigation, these exposures were assessed to be low risk overall for the school community. Classroom contacts are being advised to monitor for symptoms and seek testing if symptoms develop, effective today.

Please note: the school is not permitted to release the name or any identifying information about the confirmed cases of COVID-19 to any member of the school, community or media.

If you have not been contacted via a separate letter, your child(ren) have not been identified as being exposed to COVID-19 from these cases. However, if your child develops a new onset or worsening of any one symptom listed in Column A or any two or more symptoms listed in Column B (see table below), even if they are mild, immediately isolate them from others. It is recommended your child get tested for COVID-19 as soon as symptoms appear. You may go directly to the nearest COVID-19 testing location during their open hours. For the testing site nearest you, please visit gov.mb.ca/covid19/locations.html, or call Health Links-Info Santé at 204-788-8200 or toll-free at 1-888-315-9257.

A	В
Fever/Chills	Runny nose
Cough	Muscle aches
Sore throat/hoarse voice	Fatigue
Shortness of breath/difficulty breathing	Pink eye (conjunctivitis)
Loss of taste or smell	Headache

Vomiting or diarrhea for more than 24 hours	Skin rash of unknown cause
	Nausea or loss of appetite
	Poor feeding (if an infant)

Please note: individuals with symptoms outlined in columns A and B, who choose not to be tested, must isolate for a minimum of 10 days following the start of symptoms. Isolation can be stopped at day 10 if the individual has been symptom-free for 24 hours.

For more public health information about COVID-19, please visit <a href="https://manitoba.ca/covid19/updates/resources.html">https://manitoba.ca/covid19/updates/resources.html</a>.

Sincerely,

Dr. Lisa Richards MD MSc FRCPC

Medical Officer of Health

Winnipeg Regional Health Authority